

Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) EP 0 921 196 A1

(12) EUROPEAN PATENT APPLICATION

(43) Date of publication:  
09.06.1999 Bulletin 1999/23

(51) Int. Cl.<sup>6</sup>: C12N 15/31, C12N 9/12,  
C12Q 1/68

(21) Application number: 97121151.1

(22) Date of filing: 02.12.1997

(84) Designated Contracting States:  
AT BE CH DE DK ES FI FR GB GR IE IT LI LU MC  
NL PT SE  
Designated Extension States:  
AL LT LV MK RO SI

- Markau, Ursula  
D-82398 Polling (DE)
- Ebenbichler, Christine  
D-82387 Antdorf (DE)
- Achhammer, Gunthar  
D-82377 Penzberg (DE)

(71) Applicant:  
Roche Diagnostics GmbH  
68298 Mannheim (DE)

Remarks:

A request for correction of page 17 of the description has been filed pursuant to Rule 88 EPC. A decision on the request will be taken during the proceedings before the Examining Division (Guidelines for Examination in the EPO, A-V, 3.).

(72) Inventors:  
• Ankenbauer, Waltraud, Dr.  
D-82377 Penzberg (DE)

(54) **Modified DNA-polymerase from carboxydotherrnus hydrogenoformans and its use for coupled reverse transcription and polymerase chain reaction**

(57) A purified DNA polymerase exhibiting reverse transcriptase activity in the presence of magnesium ions and/or manganese ions having reduced or no 5'-3'-exonuclease activity and substantially no RNaseH activity and obtainable from *Carboxydotherrnus hydrogenoformans*.

EP 0 921 196 A1

## Description

[0001] The present invention relates to a modified DNA-polymerase having reverse transcriptase activity and reduced 5'-3' exonuclease activity derived from a native polymerase which is obtainable from *Carboxydotherrhus hydrogenofornans*.

Furthermore the invention relates to the field of molecular biology and provides methods for amplifying a DNA segment from an RNA template using an enzyme with reverse transcriptase activity (RT-PCR). In another aspect, the invention provides a kit for Coupled High Temperature Reverse Transcription and Polymerase Chain Reaction.

[0002] Heat stable DNA polymerases (EC 2.7.7.7. DNA nucleotidyltransferase, DNA-directed) have been isolated from numerous thermophilic organisms (for example: Kaledin et al. (1980), *Biokhimiya* 45, 644-651; Kaledin et al. (1981) *Biokhimiya* 46, 1576-1584; Kaledin et al. (1982) *Biokhimiya* 47, 1785-1791; Ruttimann et al. (1985) *Eur. J. Biochem.* 149, 41-46; Neuner et al. (1990) *Arch. Microbiol.* 153, 205-207).

For some organisms, the polymerase gene has been cloned and expressed (Lawyer et al. (1989) *J. Biol. Chem.* 264, 6427-6437; Engelke et al. (1990) *Anal. Biochem.* 191, 396-400; Lundberg et al. (1991) *Gene* 108, 1-6; Perler et al. (1992) *Proc. Natl. Acad. Sci. USA* 89, 5577-5581).

[0003] Thermophilic DNA polymerases are increasingly becoming important tools for use in molecular biology and there is growing interest in finding new polymerases which have more suitable properties and activities for use in diagnostic detection of RNA and DNA, gene cloning and DNA sequencing. At present, the thermophilic DNA polymerases mostly used for these purposes are from *Thermus* species like Taq polymerase from *T. aquaticus* (Brock et al. (1969) *J. Bacteriol.* 98, 289-297).

[0004] The term "reverse transcriptase" describes a class of polymerases characterized as RNA-dependent DNA-polymerases. All known reverse transcriptases require a primer to synthesize a DNA-transcript from an RNA template. Historically, reverse transcriptase has been used primarily to transcribe mRNA into cDNA which can then be cloned into a vector for further manipulation.

[0005] Reverse transcription is commonly performed with viral reverse transcriptases like the enzymes isolated from Avian myeloblastosis virus or Moloney murine leukemia virus. Both enzymes mentioned are active in the presence of magnesium ions but have the disadvantages to possess RNase H-activity, which destroys the template RNA during the reverse transcription reaction and have a temperature optimum at 42°C or 37 °C, respectively. Avian myeloblastosis virus (AMV) reverse transcriptase was the first widely used RNA-dependent DNA-polymerase (Verma (1977) *Biochem. Biophys. Acta* 473, 1). The enzyme has 5'-3' RNA-directed DNA polymerase activity, 5'-3' DNA directed DNA polymerase activity, and RNaseH activity. RNaseH is a processive 5'-3' ribonuclease specific for the RNA strand of RNA-DNA hybrids (Perbal (1984), *A Practical Guide to Molecular Cloning*, Wiley & Sons New York). Errors in transcription cannot be corrected because known viral reverse transcriptases lack the 3'-5' exonuclease activity necessary for proofreading (Saunders and Saunders (1987) *Microbial Genetics Applied to Biotechnology*, Croom Helm, London). A detailed study of the activity of AMV reverse transcriptase and its associated PNaseH activity has been presented by Berger et al., (1983) *Biochemistry* 22, 2365-2372.

[0006] DNA polymerases isolated from mesophilic microorganisms such as *E. coli* have been extensively characterized (see, for example, Bessmann et al. (1957) *J. Biol. Chem.* 233, 171-177 and Buttin and Kornberg (1966) *J. Biol. Chem.* 241, 5419-5427). *E. coli* DNA polymerase I (Pol I) is useful for a number of applications including: nick-translation reactions, DNA sequencing, in vitro mutagenesis, second strand cDNA synthesis, polymerase chain reactions (PCR), and blunt end formation for linker ligation (Maniatis et al., (1982) *Molecular Cloning: A Laboratory Manual* Cold Spring Harbor, New York).

[0007] Several laboratories have shown that some polymerases are capable of *in vitro* reverse transcription of RNA (Karkas (1973) *Proc. Nat. Acad. Sci. USA* 70, 3834-3838; Gulati et al. (1974) *Proc. Nat. Acad. Sci. USA* 71, 1035-1039; and Wittig and Wittig, (1978) *Nuc. Acids Res.* 5, 1165-1178). Gulati et al. found that *E. coli* Pol I could be used to transcribe Q $\beta$  viral RNA using oligo(dT)<sub>10</sub> as a primer. Wittig and Wittig have shown that *E. coli* Pol I can be used to reverse transcribe tRNA that has been enzymatically elongated with oligo(dA). However, as Gulati et al. demonstrated, the amount of enzyme required and the small size of cDNA product suggest that the reverse transcriptase activity of *E. coli* Pol I has little practical value.

[0008] Alternative methods are described using the reverse transcriptase activity of DNA polymerases of thermophilic organisms which are active at higher temperatures. Reverse transcription at higher temperatures is of advantage to overcome secondary structures of the RNA template which could result in premature termination of products. Thermostable DNA polymerases with reverse transcriptase activities are commonly isolated from *Thermus* species. These DNA polymerases however, show reverse transcriptase activity only in the presence of manganese ions. These reaction conditions are suboptimal, because in the presence of manganese ions the polymerase copies the template RNA with low fidelity.

[0009] Another feature of the commonly used reverse transcriptases is that they do not contain 3'-5' exonuclease activity. Therefore, misincorporated nucleotides cannot be removed and thus the cDNA copies from the template RNA

may contain a significant degree of mutations.

[0010] One of the known DNA polymerases having high reverse transcriptase activity is obtainable from *Thermus thermophilus* (Tth polymerase) (WO 91/0994). Tth polymerase, as well as Taq polymerase, lacks 3' to 5' exonucleolytic proofreading activity. This 3' to 5' exonuclease activity is generally considered to be desirable because it allows removal of misincorporated or unmatched bases in the newly synthesized nucleic acid sequences. Another thermophilic pol I-type DNA polymerase isolated from *Thermotoga maritima* (Tma pol) has 3' to 5' exonuclease activity. U.S. patent 5,624,833 provides means for isolating and producing Tma polymerase.

[0011] However, both DNA polymerases, Tth as well as Tma polymerase, show reverse transcriptase activity only in the presence of manganese ions.

[0012] The DNA polymerase of *Carboxydotherrnus hydrogenoformans* shows reverse transcription activity in the presence of magnesium ions and in the substantial absence of manganese ions and can be used to reverse transcribe RNA, to detect and amplify (in combination with a thermostable DNA polymerase like Taq) specific sequences of RNA. Using DNA polymerase of *Carboxydotherrnus hydrogenoformans* polymerase a high specificity of transcription is observed with short incubation times. A high specificity is observed using e.g. 5 min of incubation time and 33 units of DNA polymerase protein. With longer incubation times also with lower amounts of *Carboxydotherrnus hydrogenoformans* polymerase specific products can be obtained. However an unspecific smear of products is occurring. These unspecific products might be caused by the 5'-3' exonuclease activity of the polymerase which enables the enzyme to cleave the template at secondary structures ("RNaseH"-activity) and to create additional primers which can be elongated by the DNA polymerase activity. The thermostable DNA polymerase from *Carboxydotherrnus hydrogenoformans* has been identified and cloned and is described in the copending European application with the Application No. 96115873.0, filed October 03, 1996, and incorporated herein by reference.

[0013] In summary, reverse transcriptases as MoMULV-RT or AMV-RT perform reverse transcription in the presence of magnesium-ions. However, these enzymes act at temperatures between 37°C and 55°C. Reverse transcription at higher temperatures would be desirable because secondary structures can be overcome in the template in order to avoid premature termination of the reaction and to assure the production of cDNA without deletions.

Other enzymes e.g. DNA polymerase obtainable from *Thermus spec.* act as reverse transcriptase at temperatures up to 70°C in the presence of manganese ions. These reaction conditions are suboptimal, because in the presence of manganese ions the polymerase copies the template RNA with low fidelity and the RNA strand will be degraded. Degradation of the RNA strand occurs faster in the presence of manganese ions as in the presence of magnesium ions. Therefore, if manganese ions are present complexation of the manganese ions (e.g. with EDTA) is required after cDNA synthesis in order to obtain a higher fidelity during cDNA amplification in the subsequent PCR reaction.

[0014] Therefore, it is desirable to develop a reverse transcriptase

- which acts at higher temperatures to overcome secondary structures in the template to avoid premature termination of the reaction and to assure the production of cDNA without deletions
- which is active in the presence of magnesium ions in order to prepare cDNA from RNA templates with higher fidelity and
- which has 3'-5'-exonuclease in order to remove misincorporated nucleotides before continuation of DNA synthesis and to produce products with low mutation frequency
- which has a high specificity and produces exclusively or predominantly RT-PCR products derived from specific primer binding.

[0015] The present invention addresses these needs and provides a DNA polymerase mutant active at higher temperatures which has reverse transcriptase activity in the presence of magnesium ions and which has 3'-5' exonuclease activity and reduced or no 5'-3' exonuclease activity.

[0016] It is an object of this invention to provide a polymerase enzyme (EC 2.7.7.7.), characterized in that it has reverse transcriptase activity in the presence of magnesium ions as well as in the presence of manganese ions. In a further aspect the invention comprises a DNA polymerase having 3'-5'-exonuclease activity and reduced 5'-3' exonuclease activity. The enzyme according to the invention can be obtained from a polymerase obtainable from *Carboxydotherrnus hydrogenoformans* (Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig, DSM No. 8979). In a further aspect the invention is directed to a DNA polymerase with reduced 5'-3' exonuclease activity having reverse transcriptase activity in the presence of magnesium ions and in the substantial absence of manganese ions. In a further aspect the invention comprises a DNA polymerase having a molecular mass of about 64 to 71 kDa as determined by SDS PAGE analysis. The mutant polymerase enzyme with reduced 5'-3' exonuclease activity derived from a polymerase obtainable from *Carboxydotherrnus hydrogenoformans* is called hereinafter  $\Delta$  Chy Polymerase. In a further aspect the invention comprises a recombinant DNA sequence that encodes DNA polymerase activity of the  $\Delta$  Chy Polymerase. In a related aspect, the DNA sequence is depicted as SEQ ID No. 10 (Figure 1). In a second related aspect the invention comprises a recombinant DNA sequence that encodes

essentially amino acid residues 1 to 607 (SEQ ID No. 11, Figure 1). In a further aspect the invention comprises a recombinant DNA plasmid that comprises the DNA sequence of the invention inserted into plasmid vectors and which can be used to drive the expression of the  $\Delta$  Chy DNA polymerase in a host cell transformed with the plasmid. In a further aspect the invention includes a recombinant strain comprising the vector pDS56 carrying the  $\Delta$  Chy DNA polymerase gene and designated p $\Delta_{2-225}$ AR<sub>4</sub>. The *E. coli* strain XL1 carrying the plasmid p $\Delta_{2-225}$ AR<sub>4</sub> was deposited on the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH, Mascheroder Weg 1b, D-38124 Braunschweig DSM No. 11854 (BMTU 7307) is designated *E. coli* GA1.

[0017] In referring to a peptide chain as being comprised of a series of amino acids "substantially or effectively" in accordance with a list offering no alternatives within itself, we include within that reference any versions of the peptide chain bearing substitutions made to one or more amino acids in such a way that the overall structure and the overall function of the protein composed of that peptide chain is substantially the same as - or undetectably different to - that of the unsubstituted version. For example it is generally possible to exchange alanine and valine without greatly changing the properties of the protein, especially if the changed site or sites are at positions not critical to the morphology of the folded protein.

[0018] 3'-5' exonuclease activity is commonly referred as "proofreading" or "editing" activity of DNA polymerases. It is located in the small domain of the large fragment of Type A polymerases. This activity removes mispaired nucleotides from the 3' end of the primer terminus of DNA in the absence of nucleoside triphosphates (Kornberg A. and Baker T. A. (1992) DNA Replication W. H. Freeman & Company, New York). This nuclease action is suppressed by deoxynucleoside triphosphates if they match to the template and can be incorporated into the polymer.

[0019] The 3'-5' exonuclease activity of the claimed DNA polymerase can be measured as degradation or shortening of a 5'-digoxigenin-labeled oligonucleotide annealed to template DNA in the absence or presence of deoxyribonucleoside triphosphates or on DNA fragments in the absence or presence of deoxyribonucleoside triphosphates.

[0020] *Carboxydothermus hydrogenoformans* DNA polymerase is the first DNA polymerase isolated from thermophilic eubacteria with a higher activity in the presence of magnesium ions than in the presence of manganese ions as shown in figure 2. The reverse transcriptase activity in dependence of magnesium is of advantage since the DNA polymerases synthesize DNA with higher fidelity in the presence of magnesium than in the presence of manganese (Beckmann R. A. et al. (1985) *Biochemistry* 24, 5810-5817; Ricchetti M. and Buc H. (1993) *EMBO J.* 12, 387-396). Low fidelity DNA synthesis is likely to lead to mutated copies of the original template. In addition, Mn<sup>2+</sup> ions have been implicated in an increased rate of RNA degradation, particularly at higher temperatures and this can cause the synthesis of shortened products in the reverse transcription reaction.

[0021] The DNA sequence (SEQ ID No.: 10) of  $\Delta$  Chy polymerase and the derived amino acid sequence (SEQ ID No.: 11) of the enzyme are shown in figure 1. The molecular weight deduced from the sequence is 70,3 kDa, in SDS polyacrylamide gel electrophoresis however  $\Delta$  Chy polymerase has an electrophoretic mobility of approx. 65 kDa.

[0022] The  $\Delta$  Chy DNA Polymerase has reduced 5'-3' - exonuclease activity and has a temperature optimum at 72°C and exhibits reverse transcriptase activity at temperatures between 50 °C and 75 °C.

[0023] When using  $\Delta$  Chy DNA Polymerase obtainable from *Carboxydothermus hydrogenoformans* having reduced 5'-3' - exonuclease activity in RT-PCR as reverse transcriptase with subsequent PCR reaction using Taq-polymerase as PCR enzyme a remarkable high sensitivity is achieved (Figure 3). The sensitivity of  $\Delta$  Chy DNA Polymerase in RT-PCR is higher than the sensitivity of e.g. DNA polymerase from *Thermus thermophilus* (Tth polymerase) (Example 3, Figure 4).  $\Delta$  Chy DNA Polymerase also exhibits high sensitivity by amplifying a 1.83 kB fragment from total RNA from human muscle (Figure 5). The error rate of  $\Delta$  Chy DNA Polymerase is  $1,58 \times 10^{-4}$  mutations per nucleotide per cycle and is therewith lower than the error rate of Tth Polymerase which is  $2,37 \times 10^{-4}$  mutations per nucleotide per cycle. This results in higher fidelity of  $\Delta$  Chy DNA polymerase in comparison to Tth Polymerase.

[0024] *Carboxydothermus hydrogenoformans* was isolated from a hot spring in Kamchatka by V. Svetlichny. A sample of *C. hydrogenoformans* was deposited on the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) under the terms of the Budapest Treaty and received Accession Number DSM 8979. The thermostable polymerase isolated from *Carboxydothermus hydrogenoformans* has a molecular weight of 100 to 105 kDa. The thermostable enzyme possesses 5'-3' polymerase activity, a 3'-5' exonuclease activity and a reverse transcriptase-activity which is Mg<sup>++</sup>-dependent. The thermostable enzyme may be native or recombinant and may be used for first- and second-strand cDNA synthesis, in cDNA cloning, DNA sequencing, DNA labeling and DNA amplification.

[0025] For recovering the native protein *C. hydrogenoformans* may be grown using any suitable technique, such as the technique described by Svetlichny et al. (1991) *System. Appl. Microbiol.* 14, 205-208. After cell growth one preferred method for isolation and purification of the enzyme is accomplished using the multi-step process as follows:

[0026] The cells are thawed, suspended in buffer A (40 mM Tris-HCl, pH 7.5, 0.1 mM EDTA, 7 mM 2-mercaptoethanol, 0.4 M NaCl, 10 mM Pefabloc) and lysed by twofold passage through a Gaulin homogenizer. The raw extract is cleared by centrifugation, the supernatant dialyzed against buffer B (40 mM Tris-HCl, pH 7.5, 0.1 mM EDTA, 7 mM 2-mercaptoethanol, 10 % Glycerol) and brought onto a column filled with Heparin-Sepharose (Pharmacia). In each case the columns are equilibrated with the starting solvent and after the application of the sample washed with the threefold of its

volume with this solvent. Elution of the first column is performed with a linear gradient of 0 to 0.5 M NaCl in Buffer B. The fractions showing polymerase activity are pooled and ammonium sulfate is added to a final concentration of 20 %. This solution is applied to a hydrophobic column containing Butyl-TSK-Toyopearl (TosoHaas). The column is eluted with a falling gradient of 20 to 0 % ammonium sulfate. The pool containing the activity is dialysed and again transferred to a column of DEAE-Sepharose (Pharmacia) and eluted with a linear gradient of 0-0.5 M NaCl in buffer B. The fourth column contains Tris-Acryl-Blue (Biosepra) and is eluted as in the preceding case. Finally the active fractions are dialyzed against buffer C (20 mM Tris-HCl, pH 7.5, 0.1 mM EDTA, 7.0 mM 2-mercaptoethanol, 100 mM NaCl, 50 % Glycerol).

[0027] DNA polymerase activity was measured by incorporation of digoxigenin-labeled dUTP into the synthesized DNA and detection and quantification of the incorporated digoxigenin essentially according to the method described in Hölte, H.-J.; Sagner, G.; Kessler, C. and Schmitz, G. (1992) *Biotechniques* 12, 104-113. The reaction is performed in a reaction volume of 50 µl containing 1 or 2 µl of diluted (0.05 U - 0.01 U) DNA polymerase and 50 mM Tris-HCl, pH 8.5; 12.5 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>; 10 mM KCl; 5 mM MgCl<sub>2</sub>; 10 mM 2-mercaptoethanol; 33 µM dNTPs; 200 µg/ml BSA; 12 µg of DNase I-activated DNA from calf thymus and 0.036 µM digoxigenin-dUTP.

[0028] The samples are incubated for 30 min. at 72°C, the reaction is stopped by addition of 2 µl 0.5 M EDTA, and the tubes placed on ice. After addition of 8 µl 5 M NaCl and 150 µl of Ethanol (precooled to -20°C) the DNA is precipitated by incubation for 15 min. on ice and pelleted by centrifugation for 10 min at 13000 x rpm and 4°C. The pellet is washed with 100 µl of 70% Ethanol (precooled to -20°C) and 0.2 M NaCl, centrifuged again and dried under vacuum.

[0029] The pellets are dissolved in 50 µl Tris-EDTA (10 mM/0.1 mM; pH 7.5). 5 µl of the sample are spotted into a well of a nylon membrane bottomed white microwave plate (Pall Filtrationstechnik GmbH, Dreieich, FRG, product no: SM045BWP). The DNA is fixed to the membrane by baking for 10 min. at 70°C. The DNA loaded wells are filled with 100 µl of 0.45 µm-filtrated 1 % blocking solution (100 mM maleic acid, 150 mM NaCl, 1 % (w/v) casein, pH 7.5). All following incubation steps are done at room temperature. After incubation for 2 min. the solution is sucked through the membrane with a suitable vacuum manifold at -0.4 bar. After repeating the washing step, the wells are filled with 100 µl of a 1:10 000-dilution of Anti-digoxigenin-AP, Fab fragments (Boehringer Mannheim, FRG, no: 1093274) diluted in the above blocking solution. After incubation for 2 min. and sucking this step is repeated once. The wells are washed twice under vacuum with 200 µl each time washing-buffer 1 (100 mM maleic-acid, 150 mM NaCl, 0.3 % (v/v) Tween™ 20, pH 7.5). After washing another two times under vacuum with 200 µl each time washing-buffer 2 (10 mM Tris-HCl, 100 mM NaCl, 50 mM MgCl<sub>2</sub>, pH 9.5) the wells are incubated for 5 min. with 50 µl of CSPD™ (Boehringer Mannheim, no: 1655884), diluted 1:100 in washing-buffer 2, which serves as a chemiluminescent substrate for the alkaline phosphatase. The solution is sucked through the membrane and after 10 min. incubation the RLU/s (Relative Light Unit per second) are detected in a Luminometer e.g. MicroLumat LB 96 P (EG&G Berthold, Wilbad, FRG).

[0030] With a serial dilution of *Taq* DNA polymerase a reference curve is prepared from which the linear range serves as a standard for the activity determination of the DNA polymerase to be analyzed.

[0031] The Determination of reverse transcriptase activity is performed essentially as described for determination of DNA polymerase activity except that the reaction mixture consists of the following components: 1 µg of polydA-(dT)<sub>15</sub>, 33 µM of dTTP, 0.36 µM of digoxigenin-dUTP, 200 mg/ml BSA, 10 mM Tris-HCl, pH 8.5, 20 mM KCl, 5 mM MgCl<sub>2</sub>, 10 mM DTE and various amounts of DNA polymerase. The incubation temperature used is 50°C.

[0032] Isolation of recombinant DNA polymerase from *Carboxydotherrnus hydrogenoformans* may be performed with the same protocol or with other commonly used procedures.

[0033] The production of a recombinant form of *Carboxydotherrnus hydrogenoformans* DNA polymerase generally includes the following steps: chromosomal DNA from *Carboxydotherrnus hydrogenoformans* is isolated by treating the cells with detergent e.g. SDS and a proteinase e.g. Proteinase K. The solution is extracted with phenol and chloroform and the DNA purified by precipitation with ethanol. The DNA is dissolved in Tris/EDTA buffer and the gene encoding the DNA polymerase is specifically amplified by the PCR technique using two mixed oligonucleotides (primer 1 and 2). These oligonucleotides, described by SEQ ID No.: 1 and SEQ ID No.: 2, were designed on the basis of conserved regions of family A DNA polymerases as published by Braithwaite D. K. and Ito J. (1993) *Nucl. Acids Res.* 21, 787 - 802. The specifically amplified fragment is ligated into an vector, preferably the pCR™II vector (Invitrogen) and the sequence is determined by cycle-sequencing. Complete isolation of the coding region and the flanking sequences of the DNA polymerase gene can be performed by restriction fragmentation of the *Carboxydotherrnus hydrogenoformans* DNA with another restriction enzyme as in the first round of screening and by inverse PCR (Innis et al., (1990) PCR Protocols; Academic Press, Inc., 219-227). This can be accomplished with synthesized oligonucleotide primers binding at the outer DNA sequences of the gene part but in opposite orientation. These oligonucleotides described by SEQ ID Nos. 3 and 4, were designed on the basis of the sequences which were determined by sequencing of the first PCR product described above. As template DNA from *Carboxydotherrnus hydrogenoformans* is used which is cleaved by restriction digestion and circularized by contacting with T4 DNA ligase. To isolate the coding region of the entire polymerase gene, another PCR is performed using primers as shown in SEQ ID Nos. 5 and 6. The complete DNA polymerase gene is amplified directly from genomic DNA with primers suitable for introducing ends compatible with the linearized expression vector.

SEQ ID No. 1:

Primer 1: 5'-CCN AAY YTN CAR AAY ATH-3'

SEQ ID No. 2:

Primer 2: 5'-YTC RTC RTG NAC YTG-3'

SEQ ID No. 3:

Primer 3: 5'-GGG CGA AGA CGC TAT ATT CCT GAG C-3'

SEQ ID NO. 4:

Primer 4: 5'-GAA GCC TTA ATT CAA TCT GGG AAT AAT C-3'

SEQ ID NO. 5:

Primer 5: 5'-CGA ATT CAA TCC ATG GGA AAA GTA GTC CTG GTG GAT-3'

SEQ ID NO. 6:

Primer 6: 5'-CGA ATT CAA GGA TCC TTA CTT CGC TTC ATA CCA GTT-3'

[0034] The gene is operably linked to appropriate control sequences for expression in either prokaryotic or eucaryotic host/vector systems. The vector preferably encodes all functions required for transformation and maintenance in a suitable host, and may encode selectable markers and/or control sequences for polymerase expression. Active recombinant thermostable polymerase can be produced by transformed host cultures either continuously or after induction of expression. Active thermostable polymerase can be recovered either from host cells or from the culture media if the protein is secreted through the cell membrane.

[0035] The use of a plasmid as an appropriate vector has shown to be advantageously, particularly pDS56 (Stüber, D., Matile, H. and Garotta, G. (1990) Immunological Methods, Letkovcs, I. and Permis, B., eds). The plasmid carrying the *Carboxydotherrnus hydrogenoformans* DNA polymerase gene is then designated pAR4.

[0036] According to the present invention the use of the *E. coli* strain BL21 (DE3) pUBS520 (Brinkmann et al., (1989) gene 85, 109-114) has shown to be advantageously. The *E. coli* strain BL 21 (DEB) pUBS 520 transformed with the plasmid pAR4 is then designated AR96 (DSM No 11179).

[0037] The mutant  $\Delta$ Chy was obtained by deletion of an N-terminal fragment of the recombinant wild type *Carboxydotherrnus hydrogenoformans* DNA polymerase using inverse PCR (Innis et al., (1990) PCR Protocols; Academic Press, Inc., p 219-227). The reverse primer used is complementary to the cloning site of the expression vector pDS56 (Stüber, D., Matile, H. and Garotta, G. (1990) Immunological Methods, Letkovcs, I. and Pemis, B., eds.) at the Nco I restriction site (bases 120-151) and has the sequence:

SEQ ID No. 7:

Primer 7: 5'-CGG TAA ACC CAT GGT TAA TTT CTC CTC TTT AAT GAA TTC-3'.

[0038] This primer contains additional 7 bases at the 5' end to ensure a better binding of the Nco I restriction enzyme in the subsequent restriction enzyme cleavage. The second (forward) primer was complementary to bases 676-702 of the wild type gene and has the sequence:

SEQ ID No. 8:

Primer 8: 5'-CGG GAA TCC ATG GAA AAG CTT GCC GAA CAC GAA AAT TTA-3')

[0039] The forward primer also contained an additional Nco I restriction site and additional 7 bases at the 5'-end. Plasmid pDS56 DNA containing the polymerase-gene of *Carboxydotherrnus hydrogenoformans* at the Nco I/SamHI restriction sites was used as template for PCR. The PCR reaction was performed on the circular plasmid DNA pAR4. The fragment encoding the mutated *Carboxydotherrnus hydrogenoformans* DNA polymerase ( $\Delta$  Chy) and the vector DNA were amplified as linear DNA by PCR using the Expand High Fidelity PCR System (Boehringer Mannheim) according to the supplier's specifications (Fig. 7). The length of the gene encoding  $\Delta$  Chy is 1821 bp.

Amplification (Perkin Elmer GenAmp 9600 thermocycler) was carried out with the following conditions:

2 min 94 °C; (10 sec 94 °C; 30 sec 65 °C; 4 min 68 °C) x 10; (10 sec 94 °C; 30 sec 65 °C; 4 min 68 °C) + cycle elongation of 20 sec for each cycle) x 20; 7 min 72 °C;

After PCR the amplified DNA was purified using the High Pure PCR Product Purification Kit (Boehringer Mannheim) and digested with NcoI (3U /  $\mu$ g DNA) for 16 h (Boehringer Mannheim) according to the supplier's specifications.

For extraction with Phenol/Chloroform/Isoamylalcohol (24:24:1) the volume of the sample was raised to 100  $\mu$ l with TE. After extraction the DNA was precipitated by adding 1/10 volumes of 3M Sodium Acetate, pH 5.2 and 2 volumes of EtOH. The DNA was circularized using the Rapid DNA Ligation Kit (Boehringer Mannheim) according to the supplier's specification. The ligated products were introduced into *E. coli* XL1-Blue by transformation according to the procedure of Chung, C. T. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86, 2172-2175. Transformants were plated on L-agar containing 100  $\mu$ g/ml ampicillin to allow selection of recombinants. Colonies were picked and grown in L-broth containing 100

µg/ml ampicillin. Plasmid DNA was prepared with the High Pure Plasmid Isolation Kit (Boehringer Mannheim) according to the supplier's specification. The plasmids were screened for insertions by digestion with NcoI/BamHI. Strains containing the genes of interest were grown in L-broth supplemented with 100 µg/ml ampicillin and tested for the expression of DNA polymerase / reverse transcriptase activity by induction of exponentially growing culture with 1 mM IPTG and assaying the heat-treated extracts (72 °C) for DNA polymerase / reverse transcriptase activity as described above (determination of DNA polymerase activity and determination of reverse transcriptase activity).

[0040] The present invention provides improved methods for efficiently transcribing RNA and amplifying RNA or DNA. These improvements are achieved by the discovery and application of previously unknown properties of thermoactive DNA polymerases with reverse transcriptase activity.

[0041] The enzyme of this invention may be used for any purpose in which such enzyme activity is necessary or desired. In a particularly preferred embodiment, the enzyme catalyzes reverse transcription of RNA which is amplified as DNA by a second DNA polymerase present in the amplification reaction known as RT-PCR (Powell et al. (1987) *Cell* 50, 831-840). Any ribonucleic acid sequence, in purified or nonpurified form, can be utilized as the starting nucleic acid(s), provided it contains or is suspected to contain the specific nucleic acid sequence desired. The nucleic acid to be amplified can be obtained from any source, for example, from plasmids such as pBR322, from cloned RNA, from natural RNA from any source, including bacteria, yeast, viruses, organelles, and higher organisms such as plants and animals, or from preparations of nucleic acids made *in vitro*.

[0042] RNA may be extracted from blood, tissue material such as chorionic villi, or amniotic cells by a variety of techniques. See, e.g., Maniatis et al., (1982) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, New York) pp. 280-281. Thus the process may employ, for example, RNA, including messenger RNA, which RNA may be single-stranded or double-stranded. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized.

[0043] The amplification of target sequences from RNA may be performed to proof the presence of a particular sequence in the sample of nucleic acid to be analyzed or to clone a specific gene. Δ Chy DNA polymerase is very useful for these processes. Due to its 3'-5' exonuclease activity it is able to synthesize products with higher accuracy as the reverse transcriptases of the state of the art.

[0044] Δ Chy DNA polymerase may also be used to simplify and improve methods for detection of RNA target molecules in a sample. In these methods Δ Chy DNA polymerase from *Carboxydotherrnus hydrogenoformans* may catalyze: (a) reverse transcription and (b) second strand cDNA synthesis. The use of DNA polymerase from *Carboxydotherrnus hydrogenoformans* may be used to perform RNA reverse transcription and amplification of the resulting complementary DNA with enhanced specificity and with fewer steps than previous RNA cloning and diagnostic methods.

[0045] Another aspect of the invention comprises a kit for performing RT-PCR comprising Δ Chy polymerase, reaction buffers, nucleotide mixtures, and optionally a thermostable DNA polymerase for detection and amplification of RNA either in a one step reaction or for reverse transcription of the template RNA and subsequent amplification of the cDNA product.

#### Brief Description of the Drawings

#### [0046]

Figure 1 shows the nucleic acid and amino acid sequence of the "Klenow fragment" of Chy polymerase designated Δ Chy.

Fig. 2 shows the reverse transcriptase activity of Δ Chy in dependence of magnesium and manganese salt.

Figure 3 shows the reverse transcription and amplification of a 997 bp fragment of the β-Actin gene from total mouse liver RNA using Δ Chy and the Expand HiFi-System and decreasing amounts of RNA.

Figure 4 shows the reverse transcription and amplification of a 997 bp fragment of β-actin from total mouse liver RNA in comparison to Tth polymerase. Reverse transcription was either coupled with amplification ("one tube") using the Expand HiFi-System from Boehringer Mannheim, or after reverse transcription the Expand HiFi-System from Boehringer Mannheim was added to the reaction mixture for the subsequent amplification reaction ("two tube").

Figure 5 shows the reverse transcription and amplification of a 1,83 kb fragment of Dystrophin from total human muscle RNA.

Figure 7 shows the reverse transcription and amplification of a 324 bp fragment of β-actin from total mouse liver RNA with various amounts of Chy polymerase and various incubation times.

Figure 7 shows schematically the construction of the clone encoding Δ Chy from the clone encoding the wild type gene.

[0047] The following examples describe the invention in greater detail:

Example 1

Reverse transcription of a 324 bp  $\beta$ -actin fragment with Chy wild type DNA Polymerase used as Reverse Transcriptase followed by PCR with Taq-polymerase (Figure 6).

[0048] The reaction mixture (20  $\mu$ l) contained 200 ng total mouse liver RNA, 200  $\mu$ M dNTP, 10 mM Tris-HCl, pH 8.8, 5 mM DTT, 10 mM 2-mercaptoethanol, 15 mM KCl, 4.5 mM  $MgCl_2$ , 0.02 mg/ml BSA, 20 pmol of reverse primer ( $\beta$ -actin reverse: 5'-AAT TCG GAT GGC TAC GTA CAT GGC TG-3') and Chy-polymerase 33 units (lanes 1, 4, 7, 10, 13, 16), 13,2 units (lanes 2, 5, 8, 11, 14, 17) and 6,6 units (lanes 3, 6, 9, 12, 15, 18). Reactions were incubated for 5 min (lanes 1 to 6), 10 min (lanes 7 to 12) and 15 min (lanes 13 to 18) at 70°C.

20  $\mu$ l of the reverse transcription reaction was used as template for PCR (100  $\mu$ l reaction volume) with Taq-polymerase (Boehringer Mannheim) according to the supplier's specification using 20 pmol of forward and reverse primer (Primer sequence " $\beta$ -actin forward": 5'AGC TTG CTG TAT TCC CCT CCA TCG TG-3', primer sequence " $\beta$ -actin reverse": 5'-AAT TCG GAT GGC TAC GTA CAT GGC TG-3') and 200  $\mu$ M dNTP's.

Amplification was carried out using the following temperature profile:  
2 min 94°C; (10 sec 94 °C; 30 sec 60 °C; 30 sec 72 °C) x 30; 7 min 72 °C

Example 2

Construction of the vector expressing  $\Delta$  Chy

[0049] The mutant was obtained by deletion of an N-terminal fragment of recombinant wild type *Carboxydotherrhus hydrogenofornans* DNA polymerase using inverse PCR (Innis et al., (1990) PCR Protocols; Academic Press, Inc., p 219-227). The reverse primer used is complementary to the cloning site of the expression vector pDS56 (Stüber, D., Matile, H. and Garotta, G. (1990) Immunological Methods, Letkovcs, I. and Pernis, B., eds.) at the Nco I restriction site (bases 120-151) and has the sequence: 5'-CGG TAA ACC CAT GGT TAA TTT CTC CTC TTT AAT GAA TTC-3'. This primer contains additional 7 bases at the 5' end to ensure a better binding of the Nco I restriction enzyme in the subsequent restriction enzyme cleavage.. The second (forward) primer, was complementary to bases 676-702 of the wild type gene (sequence: 5'-CGG GAA TCC ATG GAA AAG CTT GCC GAA CAC GAA AAT TTA-3'). The forward primer also contained an additional Nco I restriction site and additional 7 bases at the 5'-end. Plasmid pDS56 DNA containing the polymerase-gene of *Carboxydotherrhus hydrogenofornans* at the Nco I/SamHI restriction sites was used as template for PCR. The PCR reaction was performed on circular plasmid DNA pAR4. The fragment of *Carboxydotherrhus hydrogenofornans* DNA polymerase ( $\Delta$ Chy) and the vector DNA were amplified as linear DNA by PCR using the Expand High Fidelity PCR System (Boehringer Mannheim) according to the supplier's specifications. The length of the gene encoding 1 Chy is 1821 bp.

Amplification (Perkin Elmer GeneAmp 9600 thermocycler) was carried out with the following conditions:  
2 min 94 °C; (10 sec 94°C; 30 sec 65 °C; 4 min 68 °C) x 10; (10 sec 94 °C; 30 sec 65 °C; 4 min 68 °C) + cycle elongation of 20 sec for each cycle) x 20; 7 min 72 °C;

After PCR the amplified DNA was purified using the High Pure PCR Product Purification Kit (Boehringer Mannheim) and digested with NcoI (3U /  $\mu$ g DNA) for 16 h (Boehringer Mannheim) according to the supplier's specifications.

For extraction with Phenol/Chloroform/Isoamylalcohol (24:24:1) the volume of the sample was raised to 100  $\mu$ l with TE. After extraction the DNA was precipitated by adding 1/10 volumes of 3M Sodium Acetate, pH 5.2 and 2 volumes of EtOH. The DNA was circularized using the Rapid DNA Ligation Kit (Boehringer Mannheim) according to the supplier's specification. The ligated products were introduced into E. coli XL1-Blue by transformation according to the procedure of Chung, C. T. et al. (1989) *Proc. Natl. Acad. Sci. USA* 86, 2172-2175. Transformants were plated on L-agar containing 100  $\mu$ g/ml ampicillin to allow selection of recombinants. Colonies were picked and grown in L-broth containing 100  $\mu$ g/ml ampicillin. Plasmid DNA was prepared with the High Pure Plasmid Isolation Kit (Boehringer Mannheim) according to the supplier's specification. The plasmids were screened for insertions by digestion with NcoI/BamHI. Strains containing the genes of interest were grown in L-broth supplemented with 100  $\mu$ g/ml ampicillin and tested for the expression of DNA polymerase / reverse transcriptase activity by induction of exponentially growing culture with 1 mM IPTG and assaying the heat-treated extracts (72 °C) for DNA polymerase / reverse transcriptase activity as described above (determination of DNA polymerase activity and determination of Reverse Transcriptase activity). (Figure 7)

Example 3

[0050] Reverse transcription and amplification of a 997 bp fragment of  $\beta$ -actin from total mouse liver RNA. Comparison of  $\Delta$  Chy with Tth polymerase in the reverse transcription reaction (Figure 4) either in a coupled RT-PCR reaction ("one tube") or in consecutive steps, reverse transcription, addition of polymerase and amplification of the cDNA prod-



uct of the first step.

"one tube" system:

- 5 [0051] The reactions (50  $\mu$ l) contained 10 mM Tris-HCl, pH 8.8 at 25 °C, 15 mM KCl, 2.5 mM MgCl<sub>2</sub>, 400  $\mu$ M of each dNTP, decreasing amounts of mouse total RNA (Clontech) as indicated in the figure, 300 nM of each primer, 60 units of  $\Delta$  Chy and 3.5 units of the Expand HiFi enzyme mix (Boehringer Mannheim GmbH). All reactions were incubated at 60 °C for 30 min (RT step). Amplification followed immediately with following cycle profile (Perkin Elmer GeneAmp 9600 thermocycler): 30 sec. at 94 °C; (30 sec at 94 °C, 30 sec at 60 °C, 1 min. at 68 °C) x 10; (30 sec. at 94°C, 30 sec. at 60°C, 1 min. at 68°C + cycle elongation of 5 sec. for each cycle) x 20; 7 min at 68 °C;

"two tube" system:

- 15 [0052] Reverse transcription is performed in 10 mM Tris-HCl, pH 8.8, 15 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 0.1 % Tween, 4.5 mM MgCl<sub>2</sub>, 2 % DMSO, 800  $\mu$ M dNTPs, 300 nmoles of each primer, 60 units of  $\Delta$  Chy, various amounts of total mouse muscle RNA as indicated in the figure. The reaction was performed in a volume of 25  $\mu$ l for 30 min at 60 °C.

- [0053] 5  $\mu$ l of this reaction are used for the amplification with the Expand HiFi-system from Boehringer Mannheim. Amplification was performed with 2.6 units of polymerase mixture in a reaction volume of 25  $\mu$ l. The following temperature cycling conditions were used: 30 sec. at 94°C, (30 sec. at 94°C, 30 sec at 60°C, 1 min at 68°C) x 10, (30 sec. at 94°C, 30 sec. at 60°C, 1 min at 68°C + cycle elongation for 5 sec for each cycle) X 20.

- 20 [0054] As a control reaction the same template-primer system was used for RT-PCR with Tth polymerase (Boehringer Mannheim). The reaction was set up according to the supplier's specifications for the "one step" variant.

#### Literature cited:

- 25 [0055]
- Beckmann R. A. et al. (1985) *Biochemistry* **24**, 5810-5817
- Berger et al., (1983) *Biochemistry* **22**, 2365-2372
- 30 Bessmann et al. (1957) *J. Biol. Chem.* **233**, 171-177
- Braithwaite D. K. and Ito J. (1993) *Nucl. Acids Res.* **21**, 787 - 802
- Brinkmann U. et al. (1989) *Gene* **85**, 109-114.
- Brock et al. (1969) *J. Bacteriol.* **98**, 289-297
- Buttin and Kornberg (1966) *J. Biol. Chem.* **241**, 5419-5427
- 35 Chung, C.T. et al. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 2172-2175
- Engelke et al. (1990) *Anal. Biochem.* **191**, 396-400.
- Gulati et al. (1974) *Proc. Nat Acad. Sci. USA* **71**, 1035-1039
- Höltke, H.-J.; Sagner, G; Kessler, C. and Schmitz, G. (1992) *Biotechniques* **12**, 104-113.
- Innis et al., (1990) *PCR Protocols*; Academic Press, Inc., 219-227
- 40 Kaledin et al. (1980), *Biokhimiya* **45**, 644-651.
- Kaledin et al. (1981) *Biokhimiya* **46**, 1576-1584.
- Kaledin et al. (1982) *Biokhimiya* **47**, 1785-1791.
- Karkas (1973) *Proc. Nat. Acad Sci. USA* **70**, 3834-3838
- Kornberg A. and Baker T.A.(1992) *DNA Replication* W. H. Freeman & Company, New York.
- 45 Lawyer et al. (1989) *J. Biol. Chem.* **264**, 6427-6437.
- Lundberg et al. (1991) *Gene* **108**, 1-6.
- Maniatis et al. (1982) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, New York.
- Neuner et al. (1990) *Arch. Microbiol.* **153**, 205-207.
- Powell et al. (1987) *Cell* **50**, 831-840
- 50 Perbal (1984), *A Practical Guide to Molecular Cloning*, Wiley & Sons New York
- Perler et al. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 5577-5581.
- Powell et al. (1987) *Cell* **50**, 831-840
- Ricchetti M. and Buc H. (1993) *EMBO J.* **12**, 387-396.
- Ruttimann et al. (1985) *Eur. J. Biochem.* **149**, 41-46.
- 55 Saunders and Saunders (1987) *Microbial Genetics Applied to Biotechnology*, Croom Helm, London
- Spanos A. and Hübscher U. (1983) *Methods in Enzymology* **91**, 263-277.
- Stüber D., Matile H. and Garotta G. (1990) *Immunological Methods*, Letkovcs, I and Pernis, B.,eds.
- Svetlichny et al. (1991) *System. Appl. Microbiol.*, **14**, 205-208.

Triglia T. et al. (1988) *Nucleic Acids Res.* 16, 8186.  
Verma (1977) *Biochem. Biophys. Acta* 473, 1  
Wittig and Wittig, (1978) *Nuc. Acids Res.* 5, 1165-1178

5

## SEQUENCE LISTING

10

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

15

(A) NAME: Boehringer Mannheim GmbH

(B) STREET: Sandhoferstr. 116

(C) CITY: Mannheim

(E) COUNTRY: DE

20

(F) POSTAL CODE (ZIP): 68305

(G) TELEPHONE: 06217595482

(H) TELEFAX: 06217594457

25

(ii) TITLE OF INVENTION: Modified DNA-Polymerase from  
carboxydotherrnus hydrogenformans and its use for Coupled Reverse  
Transcription and Polymerase Chain Reaction

30

(iii) NUMBER OF SEQUENCES: 12

## (iv) COMPUTER READABLE FORM:

35

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: IBM PC compatible

(C) OPERATING SYSTEM: PC-DOS/MS-DOS

40

(D) SOFTWARE: PatentIn Release #1.0, Version #1.30 (EPO)

## (2) INFORMATION FOR SEQ ID NO: 1:

45

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 18 base pairs

(B) TYPE: nucleic acid

50

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

55

## (ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

5 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

CCNAAYYTNC ARAAYATH

18

10

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

20

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

30

YTCRTCRTGN ACYTG

15

(2) INFORMATION FOR SEQ ID NO: 3:

35

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

40

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

45

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

50

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GGGCGAAGAC GCTATATTCC TGAGC

25

55

## (2) INFORMATION FOR SEQ ID NO: 4:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

GAAGCCTTAA TTCAATCTGG GAATAATC

28

## (2) INFORMATION FOR SEQ ID NO: 5:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: other nucleic acid

- (A) DESCRIPTION: /desc = "oligonucleotide"

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CGAATTCAAT CCATGGGAAA AGTAGTCCTG GTGGAT

36

## (2) INFORMATION FOR SEQ ID NO: 6:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

5

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

CGAATTCAAG GATCCTTACT TCGCTTCATA CCAGTT

36

10

(2) INFORMATION FOR SEQ ID NO: 7:

(i) SEQUENCE CHARACTERISTICS:

15

(A) LENGTH: 39 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

20

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

25

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

CGGTAAACCC ATGGTTAATT TCTCCTCTTT AATGAATTC

39

30

(2) INFORMATION FOR SEQ ID NO: 8:

(i) SEQUENCE CHARACTERISTICS:

35

(A) LENGTH: 39 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

40

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

45

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

CGGGAATCCA TGGAAAAGCT TGCCGAACAC GAAAATTTA

39

50

55

## (2) INFORMATION FOR SEQ ID NO: 9:

5 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 26 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 10 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid  
 15 (A) DESCRIPTION: /desc = "oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

20 AATTCGGATG GCTACGTACA TGGCTG 26

## (2) INFORMATION FOR SEQ ID NO: 10:

25 (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 1824 base pairs  
 (B) TYPE: nucleic acid  
 30 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: DNA (genomic)

(ix) FEATURE:  
 40 (A) NAME/KEY: CDS  
 (B) LOCATION:1..1824

45 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

ATG GAA AAG CTT GCC GAA CAC GAA AAT TTA GCA AAA ATA TCG AAA CAA 48  
 Met Glu Lys Leu Ala Glu His Glu Asn Leu Ala Lys Ile Ser Lys Gln  
 50 1 5 10 15

55

55

	CCC GGC CTC AAG GGG CTT TAT CTA AAG TAT GAC TTA CCG GTG TAT GAA	528
	Pro Gly Leu Lys Gly Leu Tyr Leu Lys Tyr Asp Leu Pro Val Tyr Glu	
5	165 170 175	
	GAT GTA TCT TTA AAC ATT AGA GGG TTG TTT TAT TTA AAA AAA GAA ATG	576
	Asp Val Ser Leu Asn Ile Arg Gly Leu Phe Tyr Leu Lys Lys Glu Met	
10	180 185 190	
	ATG AGA AAA ATC TTT GAG CAG GAG CAA GAA AGG TTA TTT TAT GAA ATA	624
	Met Arg Lys Ile Phe Glu Gln Glu Gln Glu Arg Leu Phe Tyr Glu Ile	
15	195 200 205	
	GAA CTT CCT TTA ACT CCA GTT CTT GCT CAA ATG GAG CAT ACC GGC ATT	672
	Glu Leu Pro Leu Thr Pro Val Leu Ala Gln Met Glu His Thr Gly Ile	
20	210 215 220	
	CAG GTT GAC CGG GAA GCT TTA AAA GAG ATG TCG TTA GAG CTG GGA GAG	720
	Gln Val Asp Arg Glu Ala Leu Lys Glu Met Ser Leu Glu Leu Gly Glu	
25	225 230 235 240	
	CAA ATT GAA GAG TTA ATC CGG GAA ATT TAT GTG CTG GCG GGG GAA GAG	768
	Gln Ile Glu Glu Leu Ile Arg Glu Ile Tyr Val Leu Ala Gly Glu Glu	
30	245 250 255	
	TTT AAC TTA AAC TCG CCC AGG CAG CTG GGA GTT ATT CTT TTT GAA AAA	816
	Phe Asn Leu Asn Ser Pro Arg Gln Leu Gly Val Ile Leu Phe Glu Lys	
35	260 265 270	
	CTT GGG CTG CCG GTA ATT AAA AAG ACC AAA ACG GGC TAC TCT ACC GAT	864
	Leu Gly Leu Pro Val Ile Lys Lys Thr Lys Thr Gly Tyr Ser Thr Asp	
40	275 280 285	
	GCG GAG GTT TTG GAA GAG CTC TTG CCT TTC CAC GAA ATT ATC GGC AAA	912
	Ala Glu Val Leu Glu Glu Leu Leu Pro Phe His Glu Ile Ile Gly Lys	
45	290 295 300	
	ATA TTG AAT TAC CGG CAG CTT ATG AAG TTA AAA TCC ACT TAT ACT GAC	960
50		
55		



ile Leu Asn Tyr Arg Gln Leu Met Lys Leu Lys Ser Thr Tyr Thr Asp  
 305 310 315 320

5 GGC TTA ATG CCT TTA ATA AAT GAG CGT ACC GGT AAA CTT CAC ACT ACT 1008  
 Gly Leu Met Pro Leu Ile Asn Glu Arg Thr Gly Lys Leu His Thr Thr  
 325 330 335

10 TTT AAC CAG ACC GGT ACT TTA ACC GGA CGC CTG GCG TCT TCG GAG CCC 1056  
 Phe Asn Gln Thr Gly Thr Leu Thr Gly Arg Leu Ala Ser Ser Glu Pro  
 340 345 350

15 AAT CTC CAA AAT ATT CCC ATC CGG TTG GAA CTC GGT CGG AAA TTA CGC 1104  
 Asn Leu Gln Asn Ile Pro Ile Arg Leu Glu Leu Gly Arg Lys Leu Arg  
 20 355 360 365

AAG ATG TTT ATA CCT TCA CCG GGG TAT GAT TAT ATT GTT TCG GCG GAT 1152  
 25 Lys Met Phe Ile Pro Ser Pro Gly Tyr Asp Tyr Ile Val Ser Ala Asp  
 370 375 380

TAT TCC CAG ATT GAA TTA AGG CTT CTT GCC CAT TTT TCC GAA GAG CCC 1200  
 30 Tyr Ser Gln Ile Glu Leu Arg Leu Leu Ala His Phe Ser Glu Glu Pro  
 385 390 395 400

AAG CTT ATT GAA GCT TAC CAA AAA GGG GAG GAT ATT CAC CGG AAA ACG 1248  
 35 Lys Leu Ile Glu Ala Tyr Gln Lys Gly Glu Asp Ile His Arg Lys Thr  
 405 410 415

GCC TCC GAG GTG TTC GGT GTA TCT TTG GAA GAA GTT ACT CCC GAG ATG 1296  
 40 Ala Ser Glu Val Phe Gly Val Ser Leu Glu Glu Val Thr Pro Glu Met  
 420 425 430

CGC GCT CAT GCC AAG TCG GTG AAC TTC GGC ATT GTT TAT GGC ATT AGT 1344  
 45 Arg Ala His Ala Lys Ser Val Asn Phe Gly Ile Val Tyr Gly Ile Ser  
 50 435 440 445

55

55

TTA ATC GCT GAA GTT GGT GCA GGC AAA AAC TGG TAT GAA GCG AAG TAA 1824  
 Leu Ile Ala Glu Val Gly Ala Gly Lys Asn Trp Tyr Glu Ala Lys \*

5                    595                    600                    605

(2) INFORMATION FOR SEQ ID NO: 11:

10                    (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 607 amino acids

(B) TYPE: amino acid

15                    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 11:

20                    Met Glu Lys Leu Ala Glu His Glu Asn Leu Ala Lys Ile Ser Lys Gln  
                     1                    5                    10                    15

25                    Leu Ala Thr Ile Leu Arg Glu Ile Pro Leu Glu Ile Ser Leu Glu Asp  
                     20                    25                    30

30                    Leu Lys Val Lys Glu Pro Asn Tyr Glu Glu Val Ala Lys Leu Phe Leu  
                     35                    40                    45

35                    His Leu Glu Phe Lys Ser Phe Leu Lys Glu Ile Glu Pro Lys Ile Lys  
                     50                    55                    60

40                    Lys Glu Tyr Gln Glu Gly Lys Asp Leu Val Gln Val Glu Thr Val Glu  
                     65                    70                    75                    80

45                    Thr Glu Gly Gln Ile Ala Val Val Phe Ser Asp Gly Phe Tyr Val Asp  
                     85                    90                    95

50                    Asp Gly Glu Lys Thr Lys Phe Tyr Ser Leu Asp Arg Leu Asn Glu Ile  
                     100                    105                    110

55                    Glu Glu Ile Phe Arg Asn Lys Lys Ile Ile Thr Asp Asp Ala Lys Gly  
                     115                    120                    125

Ile Tyr His Val Cys Leu Glu Lys Gly Leu Thr Phe Pro Glu Val Cys  
 130 135 140

5

Phe Asp Ala Arg Ile Ala Ala Tyr Val Leu Asn Pro Ala Asp Gln Asn  
 145 150 155 160

10

Pro Gly Leu Lys Gly Leu Tyr Leu Lys Tyr Asp Leu Pro Val Tyr Glu  
 165 170 175

15

Asp Val Ser Leu Asn Ile Arg Gly Leu Phe Tyr Leu Lys Lys Glu Met  
 180 185 190

20

Met Arg Lys Ile Phe Glu Gln Glu Gln Glu Arg Leu Phe Tyr Glu Ile  
 195 200 205

25

Glu Leu Pro Leu Thr Pro Val Leu Ala Gln Met Glu His Thr Gly Ile  
 210 215 220

30

Gln Val Asp Arg Glu Ala Leu Lys Glu Met Ser Leu Glu Leu Gly Glu  
 225 230 235 240

35

Gln Ile Glu Glu Leu Ile Arg Glu Ile Tyr Val Leu Ala Gly Glu Glu  
 245 250 255

40

Phe Asn Leu Asn Ser Pro Arg Gln Leu Gly Val Ile Leu Phe Glu Lys  
 260 265 270

45

Leu Gly Leu Pro Val Ile Lys Lys Thr Lys Thr Gly Tyr Ser Thr Asp  
 275 280 285

Ala Glu Val Leu Glu Glu Leu Leu Pro Phe His Glu Ile Ile Gly Lys  
 290 295 300

50

Ile Leu Asn Tyr Arg Gln Leu Met Lys Leu Lys Ser Thr Tyr Thr Asp  
 305 310 315 320

55

Gly Leu Met Pro Leu Ile Asn Glu Arg Thr Gly Lys Leu His Thr Thr  
 325 330 335  
 5  
 Phe Asn Gln Thr Gly Thr Leu Thr Gly Arg Leu Ala Ser Ser Glu Pro  
 340 345 350  
 10  
 Asn Leu Gln Asn Ile Pro Ile Arg Leu Glu Leu Gly Arg Lys Leu Arg  
 355 360 365  
 15  
 Lys Met Phe Ile Pro Ser Pro Gly Tyr Asp Tyr Ile Val Ser Ala Asp  
 370 375 380  
 20  
 Tyr Ser Gln Ile Glu Leu Arg Leu Leu Ala His Phe Ser Glu Glu Pro  
 385 390 395 400  
 25  
 Lys Leu Ile Glu Ala Tyr Gln Lys Gly Glu Asp Ile His Arg Lys Thr  
 405 410 415  
 30  
 Ala Ser Glu Val Phe Gly Val Ser Leu Glu Glu Val Thr Pro Glu Met  
 420 425 430  
 35  
 Arg Ala His Ala Lys Ser Val Asn Phe Gly Ile Val Tyr Gly Ile Ser  
 435 440 445  
 40  
 Asp Phe Gly Leu Gly Arg Asp Leu Lys Ile Pro Arg Glu Val Ala Gly  
 450 455 460  
 45  
 Lys Tyr Ile Lys Asn Tyr Phe Ala Asn Tyr Pro Lys Val Arg Glu Tyr  
 465 470 475 480  
 50  
 Leu Asp Glu Leu Val Arg Thr Ala Arg Glu Lys Gly Tyr Val Thr Thr  
 485 490 495  
 55  
 Leu Phe Gly Arg Arg Arg Tyr Ile Pro Glu Leu Ser Ser Lys Asn Arg  
 500 505 510

Thr Val Gln Gly Phe Gly Glu Arg Thr Ala Met Asn Thr Pro Leu Gln  
 515 520 525

Gly Ser Ala Ala Asp Ile Ile Lys Leu Ala Met Ile Asn Val Glu Lys  
 530 535 540

Glu Leu Lys Ala Arg Lys Leu Lys Ser Arg Leu Leu Leu Ser Val His  
 545 550 555 560

Asp Glu Leu Val Leu Glu Val Pro Ala Glu Glu Leu Glu Glu Val Lys  
 565 570 575

Ala Leu Val Lys Gly Val Met Glu Ser Val Val Glu Leu Lys Val Pro  
 580 585 590

Leu Ile Ala Glu Val Gly Ala Gly Lys Asn Trp Tyr Glu Ala Lys  
 595 600 605

(2) INFORMATION FOR SEQ ID NO: 12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 26 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: other nucleic acid

(A) DESCRIPTION: /desc = "oligonucleotide"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 12:

AGCTTGCTGT ATTCCCCTCC ATCGTG

26

## Claims

1. A purified DNA polymerase exhibiting reverse transcriptase activity in the presence of magnesium ions and/or manganese ions having reduced or no 5'-3'-exonuclease activity and substantially no RNaseH activity and obtainable from *Carboxydotherrnus hydrogenofornans*.

2. A DNA polymerase as claimed in claim 1 wherein said DNA polymerase exhibits reverse transcriptase activity in the substantial absence of manganese ions.
3. A DNA polymerase as claimed in claim 1 or 2 wherein said polymerase exhibits a reverse transcriptase activity which is manganese-dependent.
4. A DNA polymerase according to claims 1 - 3, wherein the magnesium dependent reverse transcriptase activity is higher than the manganese dependent reverse transcriptase activity of said polymerase.
5. A DNA polymerase as claimed in any of claims 1-4, wherein said polymerase is a mutant with reduced or no 5'-3' exonuclease activity derived from a naturally occurring polymerase possessing 5'-3' exonuclease activity.
6. A DNA polymerase as claimed in any of claims 1-5, wherein said polymerase has an apparent molecular weight between about 64 to 71 kDa as determined by SDS polyacrylamide electrophoresis.
7. A recombinant DNA polymerase as claimed in any of claims 1-6, wherein said polymerase is obtainable from *E. coli*, the strain being designated *E.coli* GA1.
8. An isolated DNA sequence coding for the polymerase as claimed in any one of claims 1-7.
9. A recombinant DNA sequence capable of encoding a DNA polymerase as claimed in any one of claims 1-7.
10. An isolated DNA sequence represented by the formula shown in SEQ ID No. 10.
11. A vector containing the isolated DNA sequence as claimed in any of claims 8-10.
12. A vector according to claim 11, wherein such vector is plasmid pDS56 carrying a deletion mutant of the *Carboxydotherrnus hydrogenoformans* DNA polymerase gene and is then designated p $\Delta_{2-225}$ AR<sub>4</sub>.
13. The vector according to claim 11 providing some or all of the following features:
  - (1) promoters or sites of initiation of transcription
  - (2) operators which could be used to turn gene expression on or off
  - (3) ribosome binding sites for improved translation
  - (4) transcription or translation termination sites
14. A microbial host transformed with the vector of claims 11-13.
15. A microbial host according to claim 14 wherein said transformant is *E. coli*, the strain being designated *E.coli* GA1.
16. A process for the preparation of DNA polymerase according to any of the claims 1-7 comprising the steps:
  - (a) culturing the natural strain *Carboxydotherrnus hydrogenoformans*
  - (b) suspending the cells of the natural strain in buffer
  - (c) disrupting the cells
  - (d) purifying the DNA polymerase by chromatographic steps including the use of one or more Sepharose-columns.
17. A process for the preparation of DNA polymerase according to any one of claims 1-7 comprising growing a recombinant *E. coli* strain transformed with a vector according to claims 11-13 and purifying and isolating the DNA polymerase.
18. A process of amplifying RNA, characterized in that a thermophilic DNA polymerase as claimed in any one of claims 1-7 is used in combination with a thermostable DNA polymerase.
19. A process for cDNA cloning and DNA sequencing, characterized in that a thermophilic DNA polymerase as claimed in any one of claims 1-7 is used.

20. A process for DNA labeling, characterized in that a thermophilic DNA polymerase as claimed in any one of claims 1-7 is used.

21. A process for reverse transcription of RNA to cDNA characterized in that a thermophilic DNA polymerase as claimed in any one of claims 1-7 is used.

22. A kit useful for RT-PCR comprising reverse transcription of RNA using a thermophilic DNA polymerase as claimed in any one of claims 1-7 and amplification of the cDNA product by a thermostable DNA polymerase either in a combined reaction (RT and PCR) or for consecutive reactions (RT and subsequently PCR).



Figure 1:

Nucleic acid (SEQ ID NO: 10) and protein (SEQ ID NO: 11) sequence of  $\Delta$  Chy DNA polymerase

```

1      ATGGAAGCTTGCCGAACACGAAAATTTAGCAAAATATCGAAACAATTAGCTACAATC
1      M E K L A E H E N L A K I S K Q L A T I

61     CTGCGGGAAATACCGTTAGAAATCTCCCTGGAAGATTTAAAAGTTAAAGAACCTAATTAT
21     L R E I P L E I S L E D L K V K E P N Y

121    GAAGAAGTTGCTAAATTATTTCTTCACCTTGAGTTTAAAAGCTTTTTAAAAGAAATAGAA
41     E E V A K L F L H L E F K S F L K E I E

181    CCAAAATAAAGAAAGAATACCAGGAAGGTAAAGATTTGGTGCAAGTTGAACTGTAGAA
61     P K I K K E Y Q E G K D L V Q V E T V E

241    ACGGAAGGACAGATTGCAGTAGTTTTTAGTGATGGATTTTATGTTGATGACGGGGAAAAA
81     T E G Q I A V V F S D G F Y V D D G E K

301    ACAAAGTTTTACTCTTTAGACCGGCTGAATGAAATAGAGGAAATATTTAGGAATAAAAAA
101    T K F Y S L D R L N E I E E I F R N K K

361    ATTATTACCGACGATGCCAAAGGAATTTATCATGTCTGTTTAGAAAAAGGTCTGACTTTT
121    I I T D D A K G I Y H V C L E K G L T F

421    CCCGAAGTTTGTTTGTATGCGCGGATTGCAGCTTATGTTTTAAACCGGCCGACCAAAAT
141    P E V C F D A R I A A Y V L N P A D Q N

481    CCCGGCCTCAAGGGGCTTTATCTAAAGTATGACTTACCGGTGTATGAAGATGTATCTTTA
161    P G L K G L Y L K Y D L P V Y E D V S L

541    AACATTAGAGGGTTGTTTTATTTAAAAAAGAAATGATGAGAAAAATCTTTGAGCAGGAG
181    N I R G L F Y L K K E M M R K I F E Q E

```

601 CAAGAAAGGTTATTTTATGAAATAGAACTTCCTTTAACTCCAGTTCTTGCTCAAATGGAG  
201 Q E R L F Y E I E L P L T P V L A Q M E

661 CATACCGGCATTGAGTTGACCGGGAAGCTTTAAAAGAGATGTCGTTAGAGCTGGGAGAG  
221 H T G I Q V D R E A L K E M S L E L G E

721 CAAATTGAAGAGTTAATCCGGGAAATTTATGTGCTGGCGGGGAAGAGTTTAACTTAAAC  
241 Q I E E L I R E I Y V L A G E E F N L N

781 TCGCCCAGGCAGCTGGGAGTTATTCTTTTGAAAACTTGGGCTGCCGGTAATTAAAAAG  
261 S P R Q L G V I L F E K L G L P V I K K

841 ACCAAAACGGGCTACTCTACCGATGCGGAGGTTTTGGAAGAGCTCTTGCCTTTCCACGAA  
281 T K T G Y S T D A E V L E E L L P F H E

901 ATTATCGGCAAAATATTGAATTACCGGCAGCTTATGAAGTTAAATCCACTTATACTGAC  
301 I I G K I L N Y R Q L M K L K S T Y T D

961 GGCTTAATGCCTTTAATAAATGAGCGTACCGGTAACTTCACACTACTTTTAAACCAGACC  
321 G L M P L I N E R T G K L H T T F N Q T

1021 GGTACTTTAACCGGACGCCTGGCGTCTTCGGAGCCCAATCTCCAAAATATTCCCATCCGG  
341 G T L T G R L A S S E P N L Q N I P I R

1081 TTGGAAC TCGGTCGGAAATTACGCAAGATGTTTATACCTTCACCGGGGTATGATTATATT  
361 L E L G R K L R K M F I P S P G Y D Y I

1141 GTTTCGGCGGATTATTCCCAGATTGAATTAAGGCTTCTTGCCCATTTTTCCGAAGAGCCC  
381 V S A D Y S Q I E L R L L A H F S E E P

1201 AAGCTTATTGAAGCTTACCAAAAAGGGGAGGATATTACCGGAAAACGGCCTCCGAGGTG  
401 K L I E A Y Q K G E D I H R K T A S E V

1261 TTCGGTGTATCTTTGGAAGAAGTTACTCCCGAGATGCGCGCTCATGCCAAGTCGGTGAAC  
421 F G V S L E E V T P E M R A H A K S V N

1321 TTCGGCATTGTTTATGGCATTAGTGATTTTGGTTTAGGCAGAGACTTAAAGATTCCCCGG  
441 F G I V Y G I S D F G L G R D L K I P R

1381 GAGGTTGCCGGTAAGTACATTAAAAATTATTTGCCAACTATCCCAAAGTGCGGGAGTAT  
461 E V A G K Y I K N Y F A N Y P K V R E Y

1441 CTCGATGAACTTGTCCGTACGGCAAGAGAAAAGGGATATGTGACCACTTTATTTGGGCGA  
481 L D E L V R T A R E K G Y V T T L F G R

1501 AGACGCTATATTCTGAGCTATCTTCAAAAAACCGCACGGTTCAGGGTTTGGCGAAAGG  
501 R R Y I P E L S S K N R T V Q G F G E R

1561 ACGGCCATGAATACTCCCCTTCAGGGCTCGGCTGCCGATATTATTAAGCTTGCAATGATT  
521 T A M N T P L Q G S A A D I I K L A M I

1621 AATGTAGAAAAAGAACTTAAAGCCCGTAAGCTTAAGTCCCGGCTCCTTCTTTCGGTGCAC  
541 N V E K E L K A R K L K S R L L L S V H

1681 GATGAGTTAGTTTTAGAAGTGCCGGCGGAAGAGCTGGAAGAGGTAAAAGCGCTGGTAAAA  
561 D E L V L E V P A E E L E E V K A L V K

1741 GGGGTTATGGAGTCGGTGGTTGAACTGAAAGTGCCTTTAATCGCTGAAGTTGGTGCAGGC  
581 G V M E S V V E L K V P L I A E V G A G

1801 AAAAAGTGGTATGAAGCGAAGTAA  
601 K N W Y E A K \*

Figure 2:




Figure 3:




Figure 4:




Figure 5:

M	50	10	50	10	5	1	500	10	M
	0	0	ng	ng	ng	ng	pg	pg	
			ng	ng					




Figure 6:

1 2 3 4 5 6 7 8 9 10 11 12 M 13 14 15 16 17 18





Figure 7:





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 97 12 1151

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	WO 96 10640 A (LIFE TECHNOLOGIES INC) 11 April 1996  see the claims * abstract; examples 4-7, 12, 15 * * page 10, line 28 - page 11, line 30 * * page 14, line 21 - page 15, line 2 * ---	1-3, 5, 6, 8, 9, 11, 13, 14, 17-22	C12N15/31 C12N9/12 C12Q1/68
X	WO 92 03556 A (CETUS CORP) 5 March 1992  * abstract; examples 4-7 * * page 2, paragraph 4 - page 3, paragraph 3 * * page 12, paragraph 4 - page 17, paragraph 2 * * page 32, paragraph 2 - page 34, paragraph 2 * ---	1-3, 5, 6, 8, 9, 11, 13, 14, 17-22	
X	WO 91 09944 A (CETUS CORP) 11 July 1991  see the claims * abstract; examples 4-6, 8 * * page 16, line 3 - page 15 * ---	1-3, 5, 6, 8, 9, 11, 13, 14, 17-22	TECHNICAL FIELDS SEARCHED (Int.Cl.6) C12N C12Q
A	SVETLICHNY V A ET AL: "Carboxydotherrnus hydrogenoformans gen. nov., sp. nov., a CO-utilizing thermophilic anaerobic bacterium from hydrothermal environments of Lunashir Island" SYSTEM. APPL. MICROBIOL., vol. 14, no. 3, 1991, pages 254-260, XP002074994 * the whole document * --- -/--		
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 19 August 1998	Examiner Oderwald, H
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

EPO FORM 1503 03/82 (P04C01)





European Patent  
Office

## EUROPEAN SEARCH REPORT

Application Number  
EP 97 12 1151

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
T	EP 0 834 569 A (BOEHRINGER MANNHEIM GMBH) 8 April 1998  see the claims * abstract: figures 5,6; example 6 * -----	1-6,8,9, 11,13, 14,16-22	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
The present search report has been drawn up for all claims			
Place of search <b>THE HAGUE</b>		Date of completion of the search <b>19 August 1998</b>	Examiner <b>Oderwald, H</b>
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03/82 (P04C01)

**THIS PAGE BLANK (USPTO)**